

Deformation of the ABJM Theory

Mir Faizal

Mathematical Institute, University of Oxford
Oxford OX1 3LB, United Kingdom

March 2, 2013

Abstract

In this paper we analyse the ABJM theory on deformed spacetime. We show that this theory reduces to a deformed super-Yang-Mills theory when one of the scalar superfields is given a non-vanishing vacuum expectation value. Our analyse is done in $N = 1$ superspace formalism.

1 Introduction

The low energy action for multiple $M2$ -branes is thought to be given by the ABJM theory [1, 2]. This theory is a three dimensional Chern-Simons-matter theory with gauge group $U(N)_k \times U(N)_{-k}$ at levels k and $-k$ on the world-volume of N $M2$ -membranes placed at the fixed point of R^8/Z_k . This theory explicitly realizes $N = 6$ supersymmetry and it is expected to be enhanced to full $N = 8$ supersymmetry for $k = 1, 2$ [3]. The ABJM theory coincides with the BLG theory for the only known example of the Lie 3-algebra [4, 5, 6, 7, 8]. Both the BLG theory and the ABJM theory have been analysed in the $N = 1$ superfield formalism [9, 10] and the Higgs mechanism for both these theories has also been studied in the $N = 1$ superspace formalism [11, 12].

The spacetime noncommutativity arises in the string theory because of coupling the theory to a NS background [13, 14]. Other backgrounds fields can similarly cause other deformations of the theory. A non-anticommutative deformation of the theory is caused by a RR background [21, 22]. Furthermore, the commutator of the spacetime coordinates with the fermionic coordinates does not vanish in the presence of a gravitino background [23]. Field theories with these kind of deformations have been thoroughly studied [22]-[28]. In fact, it is well known that non-anticommutative deformation of a theory breaks half of the supersymmetry of that theory explicitly. In this paper we will study those deformations of the M -theory which do not break any supersymmetry.

As there is a duality between M -theory and II string theory, we expect that any deformation on the M -theory side will also correspond to some deformation on the string theory side. In fact, $M2$ -branes in M -theory are analogous objects to strings in string theory. This is because just like strings can end on D -brane in string theory, $M2$ -branes can end on $M5$ -branes in M -theory. Furthermore, as a three-form field strength occurs naturally in M -theory, we expect that coupling the ABJM theory to a background three-form field could lead to a deformation

of the ABJM theory, just like coupling of D -branes to a background two-form field strength leads to a noncommutative deformation of the theory.

Coupling of the ABJM theory to background fields can have interesting uses in understanding the theory of multiple $M5$ -branes. It may be noted that even though the action for a single $M5$ -brane is known, the action for multiple $M5$ -branes is not known [31]-[35]. Furthermore, in M -theory the action of a single $M5$ -brane can be obtained by analyzing the κ -symmetry of the open membrane ending on it [29]. It might be possible to perform a similar analysis using the ABJM theory coupled to a background three-form field strength and gain insight into the dynamics of multiple $M5$ -branes.

2 ABJM Theory in $N = 1$ Superspace

Now we review the classical Lagrangian density for the ABJM theory in $N = 1$ superspace formalism with the gauge group $U(N) \times U(N)$,

$$\mathcal{L}_c = \mathcal{L}_M + \mathcal{L}_{CS} - \tilde{\mathcal{L}}_{CS}, \quad (1)$$

where $\mathcal{L}_{CS}$ and $\tilde{\mathcal{L}}_{CS}$ are deformed Chern-Simons theories with gauge group's $U(N)$ from $U(N) \times U(N)$ respectively. They can thus be expressed as

$$\begin{aligned} \mathcal{L}_{CS} &= \frac{k}{2\pi} \int d^2\theta \, Tr [\Gamma^a \Omega_a]_1, \\ \tilde{\mathcal{L}}_{CS} &= \frac{k}{2\pi} \int d^2\theta \, Tr [\tilde{\Gamma}^a \tilde{\Omega}_a]_1, \end{aligned} \quad (2)$$

where k is an integer and

$$\Omega_a = \omega_a - \frac{1}{6} [\Gamma^b, \Gamma_{ab}] \quad (3)$$

$$\omega_a = \frac{1}{2} D^b D_a \Gamma_b - \frac{i}{2} [\Gamma^b, D_b \Gamma_a] - \frac{1}{6} [\Gamma^b, \{\Gamma_b, \Gamma_a\}], \quad (4)$$

$$\Gamma_{ab} = -\frac{i}{2} [D_{(a} \Gamma_{b)} - i \{\Gamma_a, \Gamma_b\}],$$

$$\tilde{\Omega}_a = \tilde{\omega}_a - \frac{1}{6} [\tilde{\Gamma}^b, \tilde{\Gamma}_{ab}] \quad (5)$$

$$\tilde{\omega}_a = \frac{1}{2} D^b D_a \tilde{\Gamma}_b - \frac{i}{2} [\tilde{\Gamma}^b, D_b \tilde{\Gamma}_a] - \frac{1}{6} [\tilde{\Gamma}^b, \{\tilde{\Gamma}_b, \tilde{\Gamma}_a\}], \quad (6)$$

$$\tilde{\Gamma}_{ab} = -\frac{i}{2} [D_{(a} \tilde{\Gamma}_{b)} - i \{\tilde{\Gamma}_a, \tilde{\Gamma}_b\}]. \quad (7)$$

Here the super-derivative D_a is given by

$$D_a = \partial_a + (\gamma^\mu \partial_\mu)_a^b \theta_b, \quad (8)$$

and $'|'$ means that the quantity is evaluated at $\theta_a = 0$. In component form the Γ_a and $\tilde{\Gamma}_a$ are given by

$$\begin{aligned} \Gamma_a &= \chi_a + B\theta_a + \frac{1}{2} (\gamma^\mu)_a A_\mu + i\theta^2 \left[\lambda_a - \frac{1}{2} (\gamma^\mu \partial_\mu \chi)_a \right], \\ \tilde{\Gamma}_a &= \tilde{\chi}_a + \tilde{B}\theta_a + \frac{1}{2} (\gamma^\mu)_a \tilde{A}_\mu + i\theta^2 \left[\tilde{\lambda}_a - \frac{1}{2} (\gamma^\mu \partial_\mu \tilde{\chi})_a \right]. \end{aligned} \quad (9)$$

Thus in component form these Lagrangian are given by

$$\begin{aligned}
\mathcal{L}_{cs} &= \frac{k}{4\pi} \left(2 \left(\epsilon^{\mu\nu\rho} A_\mu \partial_\nu A_\rho + \frac{2i}{3} A_\mu A_\nu A_\rho \right) \right. \\
&\quad \left. + E^a E_a + \mathcal{D}_\mu (\chi^a (\gamma^\mu)_a^b E_b) \right), \\
\tilde{\mathcal{L}}_{cs} &= \frac{k}{4\pi} \left(2 \left(\epsilon^{\mu\nu\rho} \tilde{A}_\mu \partial_\nu \tilde{A}_\rho + \frac{2i}{3} \tilde{A}_\mu \tilde{A}_\nu \tilde{A}_\rho \right) \right. \\
&\quad \left. + \tilde{E}^a \tilde{E}_a + \tilde{\mathcal{D}}_\mu (\tilde{\chi}^a (\gamma^\mu)_a^b \tilde{E}_b) \right).
\end{aligned} \tag{10}$$

The Lagrangian density for the matter fields is given by

$$\mathcal{L}_M = \frac{1}{4} \int d^2\theta \, Tr \left[[\nabla^a X^{I\dagger} \nabla_a X_I] + \mathcal{V} \right], \tag{11}$$

where

$$\begin{aligned}
\nabla_a X^I &= D_a X^I + i\Gamma_a X^I - iX^I \tilde{\Gamma}_a, \\
\nabla_a X^{I\dagger} &= D_a X^{I\dagger} - iX^{I\dagger} \Gamma_a + i\tilde{\Gamma}_a X^{I\dagger},
\end{aligned} \tag{12}$$

and $\mathcal{V}$ is the potential term given by

$$\mathcal{V} = \frac{16\pi}{k} \epsilon^{IJ} \epsilon_{KL} [X_I X^{K\dagger} X_J X^{L\dagger}]. \tag{13}$$

Now if the full finite gauge transformation under which this ABJM theory invariant are given by

$$\begin{aligned}
\Gamma_a &\rightarrow iu \nabla_a u^{-1}, & \tilde{\Gamma}_a &\rightarrow i\tilde{u} \nabla_a \tilde{u}^{-1}, \\
X^I &\rightarrow u X^I \tilde{u}^{-1}, & X^{I\dagger} &\rightarrow \tilde{u} X^{I\dagger} u^{-1},
\end{aligned} \tag{14}$$

where

$$\begin{aligned}
u &= [\exp(i\Lambda^A T_A)], \\
\tilde{u} &= [\exp(i\tilde{\Lambda}^A T_A)].
\end{aligned} \tag{15}$$

Thus now the infinitesimal gauge transformations for these fields is given by

$$\begin{aligned}
\delta\Gamma_a &= \nabla_a \Lambda, & \delta\tilde{\Gamma}_a &= \nabla_a \tilde{\Lambda}, \\
\delta X^I &= i(\Lambda X^I - X^I \tilde{\Lambda}), & \delta X^{I\dagger} &= i(\tilde{\Lambda} X^{I\dagger} - X^{I\dagger} \Lambda),
\end{aligned} \tag{16}$$

The Lagrangian for the ABJM theory is invariant under these gauge transformations

$$\begin{aligned}
\delta\mathcal{L}_{ABJM} &= \delta\mathcal{L}_{kcs}(\Gamma) - \delta\tilde{\mathcal{L}}_{-kcs}(\tilde{\Gamma}) + \delta\mathcal{L}_M \\
&= 0.
\end{aligned} \tag{17}$$

3 Deformed ABJM Theory

In this section we shall construct a three dimensional Chern-Simons theory on a deformed superspace. In order to analyse deformation of the superspace both

the Grassman coordinates and the spacetime coordinates are promoted to operators and a deformation of there superalgebra is imposed. In four dimensions $N = 1$ supersymmetry is generated by the supersymmetric generator Q_A which can be split into Q_a and $Q_{\dot{a}}$. Furthermore, it is possible to break the supersymmetry corresponding to $Q_{\dot{a}}$ and leave the supersymmetry corresponding to Q_a intact or vice versa [22]. Thus, it is possible to construct theories with $N = 1/2$ supersymmetry in four dimensions. However, in three dimensions both Q_a and $Q_{\dot{a}}$ act as independent supercharges. So, we can view a theory with $N = 1$ supersymmetry in four dimensions as a theory with $N = 2$ supersymmetry in three dimensions. Thus, $N = 1/2$ supersymmetry in four dimensions corresponds to $N = 1$ supersymmetry in three dimensions [30]. It is not possible to obtain a $N = 1/2$ theory in three dimensions as there are not enough degrees of freedom in three dimensions to do that. So, if we deform a theory with $N = 1$ supersymmetry in three dimensions, we can either retain all the supersymmetry or break all of it. However, we cannot partially break $N = 1$ supersymmetry in three dimensions. As supersymmetry is very important in the analysis of the ABJM theory we will deform the superspace algebra in such a way that we do not break any supersymmetry. To do so we promote θ^a and y^μ to operators $\hat{\theta}^a$ and $\hat{y}^\mu$ which satisfy the following superspace algebra,

$$[\hat{y}^\mu, \hat{y}^\nu] = B^{\mu\nu}, \quad [\hat{y}^\mu, \hat{\theta}^a] = A^{\mu a}. \quad (18)$$

We then use Weyl ordering and express the Fourier transformation of this superfield as,

$$\hat{\Gamma}_a(\hat{y}, \hat{\theta}) = \int d^4k \int d^2\pi e^{-ik\hat{y} - \pi\hat{\theta}} \Gamma_a(k, \pi). \quad (19)$$

Thus, we obtain a one to one map between a function of $\hat{\theta}, \hat{y}$ to a function of ordinary superspace coordinates θ, y via

$$\Gamma_a(y, \theta) = \int d^4k \int d^2\pi e^{-iky - \pi\theta} \Gamma_a(k, \pi). \quad (20)$$

We can express the product of two fields $\hat{\Gamma}^a(\hat{y}, \hat{\theta})\hat{\Gamma}_a(\hat{y}, \hat{\theta})$ on this deformed superspace as

$$\begin{aligned} \hat{\Gamma}^a(\hat{y}, \hat{\theta})\hat{\Gamma}_a(\hat{y}, \hat{\theta}) &= \int d^4k_1 d^4k_2 \int d^2\pi_1 d^2\pi_2 \exp -i((k_1 + k_2)\hat{y} + (\pi_1 + \pi_2)\hat{\theta}) \\ &\quad \times \exp(i\Delta)\Gamma^a(k_1, \pi_1)\Gamma_a(k_2, \pi_2), \end{aligned} \quad (21)$$

where

$$\exp(i\Delta) = \exp -\frac{i}{2} (B^{\mu\nu} k_\mu^2 k_\nu^1 + A^{\mu a} (\pi_a^2 k_\mu^1 - k_\mu^2 \pi_a^1)), \quad (22)$$

This motivates the definition of the star product between ordinary functions, which is now defined as

$$\begin{aligned} \Gamma^a(y, \theta) \star \Gamma_a(y, \theta) &= \exp -\frac{i}{2} (B^{\mu\nu} \partial_\mu^2 \partial_\nu^1 + A^{\mu a} (D_a^2 \partial_\mu^1 - \partial_\mu^2 D_a^1)) \\ &\quad \times \Gamma^a(y_1, \theta_1) \Gamma_a(y_2, \theta_2) \big|_{y_1=y_2=y, \theta_1=\theta_2=\theta}. \end{aligned} \quad (23)$$

Here we have defined the star product between ordinary functions using super-derivative D_a rather than ∂_a because they commute with the generators of the supersymmetry Q_a [30],

$$\{Q_a, D_b\} = 0. \quad (24)$$

Thus, to write the ABJM in this deformed superspace, we now use

$$\begin{aligned}\hat{\Omega}_a &= \hat{\omega}_a - \frac{1}{6}[\hat{\Gamma}^b, \hat{\Gamma}_{ab}] \\ &= \omega_a - \frac{1}{6}[\Gamma^b, \Gamma_{ab}]_\star\end{aligned}\quad (25)$$

$$\begin{aligned}\hat{\omega}_a &= \frac{1}{2}D^b D_a \hat{\Gamma}_b - \frac{i}{2}[\hat{\Gamma}^b, D_b \hat{\Gamma}_a] - \frac{1}{6}[\hat{\Gamma}^b, \{\hat{\Gamma}_b, \hat{\Gamma}_a\}] \\ &= \frac{1}{2}D^b D_a \Gamma_b - \frac{i}{2}[\Gamma^b, D_b \Gamma_a]_\star - \frac{1}{6}[\Gamma^b, \{\Gamma_b, \Gamma_a\}_\star],\end{aligned}\quad (26)$$

$$\begin{aligned}\hat{\Gamma}_{ab} &= -\frac{i}{2}[D_{(a} \hat{\Gamma}_{b)} - i\{\hat{\Gamma}_a, \hat{\Gamma}_b\}] \\ &= -\frac{i}{2}(D_{(a} \Gamma_{b)} - i\{\Gamma_a, \Gamma_b\}_\star),\end{aligned}\quad (27)$$

$$\begin{aligned}\hat{\hat{\Omega}}_a &= \hat{\hat{\omega}}_a - \frac{1}{6}[\hat{\hat{\Gamma}}^b, \hat{\hat{\Gamma}}_{ab}] \\ &= \tilde{\omega}_a - \frac{1}{6}[\tilde{\Gamma}^b, \tilde{\Gamma}_{ab}]_\star\end{aligned}\quad (28)$$

$$\begin{aligned}\hat{\hat{\omega}}_a &= \frac{1}{2}D^b D_a \hat{\hat{\Gamma}}_b - \frac{i}{2}[\hat{\hat{\Gamma}}^b, D_b \hat{\hat{\Gamma}}_a] - \frac{1}{6}[\hat{\hat{\Gamma}}^b, \{\hat{\hat{\Gamma}}_b, \hat{\hat{\Gamma}}_a\}] \\ &= \frac{1}{2}D^b D_a \tilde{\Gamma}_b - \frac{i}{2}[\tilde{\Gamma}^b, D_b \tilde{\Gamma}_a]_\star - \frac{1}{6}[\tilde{\Gamma}^b, \{\tilde{\Gamma}_b, \tilde{\Gamma}_a\}_\star],\end{aligned}\quad (29)$$

$$\begin{aligned}\tilde{\Gamma}_{ab} &= -\frac{i}{2}[D_{(a} \tilde{\Gamma}_{b)} - i\{\tilde{\Gamma}_a, \tilde{\Gamma}_b\}] \\ &= -\frac{i}{2}[D_{(a} \tilde{\Gamma}_{b)} - i\{\tilde{\Gamma}_a, \tilde{\Gamma}_b\}_\star].\end{aligned}\quad (30)$$

In order to analyse the gauge transformations of this deformed ABJM theory it will be useful to define

$$\begin{aligned}u &= [\exp(i\hat{\Lambda}^A T_A)] = [\exp(i\Lambda^A T_A)]_\star, \\ \tilde{u} &= [\exp(i\hat{\hat{\Lambda}}^A T_A)] = [\exp(i\tilde{\Lambda}^A T_A)]_\star.\end{aligned}\quad (31)$$

The star product reduces to the usual Moyal star product for the bosonic non-commutativity in the limit $C^{ab} = A^{a\mu} = 0$ and for $A^{a\mu} = C^{ab} = 0$ it reduces to the standard fermionic star product.

Now we construct the classical Lagrangian density with the gauge group $U(N) \times U(N)$, on this deformed superspace,

$$\mathcal{L}_c = \mathcal{L}_M + \mathcal{L}_{CS} - \tilde{\mathcal{L}}_{CS}, \quad (32)$$

where $\mathcal{L}_{CS}$ and $\tilde{\mathcal{L}}_{CS}$ are deformed Chern-Simons theories with gauge group's $U(N)$ from $U(N) \times U(N)$ respectively. They can thus be expressed as

$$\begin{aligned}\mathcal{L}_{CS} &= \frac{k}{2\pi} \int d^2\theta \, Tr \left[\hat{\Gamma}^a \hat{\Omega}_a \right]_{|} \\ &= \frac{k}{2\pi} \int d^2\theta \, Tr \left[\Gamma^a \star \Omega_a \right]_{|}, \\ \tilde{\mathcal{L}}_{CS} &= \frac{k}{2\pi} \int d^2\theta \, Tr \left[\hat{\hat{\Gamma}}^a \hat{\hat{\Omega}}_a \right]_{|} \\ &= \frac{k}{2\pi} \int d^2\theta \, Tr \left[\tilde{\Gamma}^a \star \tilde{\Omega}_a \right]_{|}.\end{aligned}\quad (33)$$

The Lagrangian for the matter fields is given by

$$\mathcal{L}_M = \frac{1}{4} \int d^2 \theta \text{Tr} \left[[\nabla^a \hat{X}^{I\dagger} \nabla_a \hat{X}_I] + \hat{\mathcal{V}} \right], \quad (34)$$

where

$$\begin{aligned} \nabla_a \hat{X}^I &= D_a \hat{X}^I + i \hat{\Gamma}_a \hat{X}^I - i \hat{X}^I \hat{\Gamma}_a, \\ \nabla_a \hat{X}^{I\dagger} &= D_a \hat{X}^{I\dagger} - i \hat{X}^{I\dagger} \hat{\Gamma}_a + i \hat{\Gamma}_a \hat{X}^{I\dagger}, \end{aligned} \quad (35)$$

and $\hat{\mathcal{V}}$ is the potential term given by

$$\hat{\mathcal{V}} = \frac{16\pi}{k} \epsilon^{IJ} \epsilon_{K L} [\hat{X}_I \hat{X}^{K\dagger} \hat{X}_J \hat{X}^{L\dagger}]. \quad (36)$$

Now we can also express the Lagrangian for the matter fields as

$$\mathcal{L}_M = \frac{1}{4} \int d^2 \theta \text{Tr} \left[[\nabla^a \star X^{I\dagger} \star \nabla_a \star X_I] + \mathcal{V}_\star \right], \quad (37)$$

where

$$\begin{aligned} \nabla_a \star X^I &= D_a \star X^I + i \Gamma_a \star X^I - i \star X^I \tilde{\Gamma}_a, \\ \nabla_a \star X^{I\dagger} &= D_a \star X^{I\dagger} - i \star X^{I\dagger} \tilde{\Gamma}_a + i \tilde{\Gamma}_a \star X^{I\dagger}, \end{aligned} \quad (38)$$

and $\mathcal{V}_\star$ is the potential term given by

$$\mathcal{V}_\star = \frac{16\pi}{k} \epsilon^{IJ} \epsilon_{K L} [\star X_I \star X^{K\dagger} \star X_J \star X^{L\dagger}]. \quad (39)$$

Now if the full finite gauge transformation under which this ABJM theory invariant are given by

$$\begin{aligned} \Gamma_a &\rightarrow i u \star \nabla_a \star u^{-1}, & \tilde{\Gamma}_a &\rightarrow i \tilde{u} \star \nabla_a \star \tilde{u}^{-1}, \\ X^I &\rightarrow u \star X^I \star \tilde{u}^{-1}, & X^{I\dagger} &\rightarrow \tilde{u} \star X^{I\dagger} \star u^{-1}. \end{aligned} \quad (40)$$

Thus now the infinitesimal gauge transformations for these fields is given by

$$\begin{aligned} \delta \Gamma_a &= \nabla_a \star \Lambda, & \delta \tilde{\Gamma}_a &= \nabla_a \star \tilde{\Lambda}, \\ \delta X^I &= i(\Lambda \star X^I - X^I \star \tilde{\Lambda}), & \delta X^{I\dagger} &= i(\tilde{\Lambda} \star X^{I\dagger} - X^{I\dagger} \star \Lambda), \end{aligned} \quad (41)$$

The Lagrangian for the ABJM theory is invariant under these gauge transformations

$$\begin{aligned} \delta \mathcal{L}_{ABJM} &= \delta \mathcal{L}_{kcs}(\Gamma) - \delta \tilde{\mathcal{L}}_{-kcs}(\tilde{\Gamma}) + \delta \mathcal{L}_M \\ &= 0. \end{aligned} \quad (42)$$

4 Higgs Mechanism

Now we take the vacuum expectation value of one of the scalar superfields say X , to be a non-zero,

$$\langle X \rangle = \nu \neq 0. \quad (43)$$

This spontaneously breaks the symmetry from $U(N) \times U(N)$ to its diagonal subgroup, $U(N)$. Now let A_a be superfield associated with the broken gauge and B_a be associated with the unbroken gauge group. Then we have

$$\begin{aligned} A_a &= \frac{1}{2} (\Gamma_a - \tilde{\Gamma}_a), \\ B_a &= \frac{1}{2} (\Gamma_a + \tilde{\Gamma}_a), \end{aligned} \quad (44)$$

Now we can write the Chern-Simons part of the Lagrangian as

$$\mathcal{L}_{CS} = \frac{k}{2\pi} \int d^2\theta \, Tr \left[A^a \star \left[W_a + \frac{1}{6} [A^b, A_{ab}]_\star \right] \right]_|, \quad (45)$$

where

$$\begin{aligned} W_a &= \frac{1}{2} D^b D_a B_b - \frac{i}{2} [B^b, D_b B_a]_\star - \frac{1}{6} [B^b, \{B_b, B_a\}_\star]_\star, \\ A_{ab} &= -\frac{i}{2} [D_a B_b] - i \{A_a, B_b\}_\star. \end{aligned} \quad (46)$$

Now as the gauge group is broken down to its diagonal subgroup, we can integrate out the field A_a by using its equations of motion. Using this value of A_a thus obtained we get the a theory whose first term corresponds to noncommutative super-Yang-Mills theory. Thus the kinetic part of this theory has the terms

$$\mathcal{L}_{YM} = \frac{k^2}{16\pi^2\nu^2} \int d^2\theta \, [W^a \star W_a + [\mathcal{O}]_\star]_|. \quad (47)$$

There are higher derivative terms in this action corresponding to different whose origin is not the noncommutative nature of the theory we have analysed. Similarly in the kinetic part of the matter field also contains higher derivatives. However, the first term is a usual gauge theory term with the covariant derivatives corresponding to the unbroken gauge field B_a .

$$\mathcal{L}_{KM} = \frac{k^2}{16\pi^2\nu^2} \int d^2\theta \, [\nabla^a \star X^{I\dagger} \star \nabla_a \star X_I + [\mathcal{O}]_\star]_|, \quad (48)$$

where

$$\begin{aligned} \nabla_a \star X^I &= D_a X^I - i B_a \star X^I, \\ \nabla_a \star X^{I\dagger} &= D_a X^{I\dagger} + i B_a \star X^{I\dagger}. \end{aligned} \quad (49)$$

The potential term can be now rewritten as $V_\star$ using equation of motion of A_a . We identify the Yang-Mills coupling with

$$g_{YM} = \frac{2\pi\nu}{k}. \quad (50)$$

It is possible to keep g_{YM} fixed in the limit $\nu \rightarrow \infty$ and $k \rightarrow \infty$. So we will now only consider the leading order terms in powers of ν and k . thus we can write the full Lagrangian as

$$\mathcal{L}_T = \frac{1}{g_{YM}^2} \int d^2\theta \, [W^a \star W_a + \nabla^a \star X^{I\dagger} \star \nabla_a \star X_I + V_\star]_| \quad (51)$$

Now if the full finite gauge transformation under which this theory is invariant are given by

$$\begin{aligned} B_a &\rightarrow iv\star\nabla_a\star v^{-1}, & X^I &\rightarrow v\star X^I, \\ X^{I\dagger} &\rightarrow X^{I\dagger}\star v^{-1}, \end{aligned} \quad (52)$$

where

$$v = u + \tilde{u}. \quad (53)$$

We can write now v as follows

$$v = [\exp(i\lambda^A T_A)]_\star. \quad (54)$$

It may also be noted that

$$\delta W_a = v\star W_a\star v^{-1}, \quad (55)$$

Now the infinitesimal gauge transformations for these fields can be written as

$$\begin{aligned} \delta B_a &= \nabla_a\star\lambda, & \delta X^I &= i\lambda\star X^I, \\ \delta X^{I\dagger} &= -iX^{I\dagger}\star\lambda. \end{aligned} \quad (56)$$

The Lagrangian for the ABJM theory is invariant under these gauge transformations

$$\delta\mathcal{L}_T = 0. \quad (57)$$

Thus this theory is just a super-Yang-Mills theory deformed by the background fields. This deformed super-Yang-Mills theory arises as a result of coupling D -branes to background fields. This motivates the question if it is possible to write a non-linear Born-Infidel type extension to ABJM theory. As the gauge part is purely topological we expect no change in that part. However, it might be possible to write the kinetic part of the matter fields as a Born-Infidel type theory.

5 Conclusion

In this paper we have analysed the ABJM theory in $N = 1$ superspace on non-commutative spacetime. It was shown that this theory is invariant under noncommutative gauge transformations, which in commutative limit reduce to regular gauge transformations for the ABJM theory. Furthermore, it was demonstrated that this theory reduces to the noncommutative super-Yang-Mills theory when one of the scalar fields is given an vacuum expectation value.

Recently supersymmetric Chern-Simons theory has also been used to study fractional quantum Hall effect via holography [36] and analyse the AdS_4/CFT_3 correspondence [37]-[41]. It will be interesting to analyse similar effects for this deformed ABJM theory. It will also be interesting to analyse the BRST and the anti-BRST symmetries of this model. These symmetries for noncommutative deformation of the ABJM theory in various cases have been already analysed [10]. It will be interesting to analyse the BRST and the anti-BRST symmetries of the ABJM theory with non-anticommutative deformation.

References

- [1] I. L. Buchbinder, E.A. Ivanov, O. Lechtenfeld, N. G. Pletnev, I. B. Samsonov and B. M. Zupnik, JHEP. 0903, 096 (2009)
- [2] O. Aharony, O. Bergman, D. L. Jafferis and J. Maldacena, JHEP. 0810, 091 (2008)
- [3] O-Kab Kwon, P. Oh and J. Sohn, JHEP. 0908, 093 (2009)
- [4] J. Bagger and N. Lambert, Phys. Rev. D75, 045020 (2007)
- [5] J. Bagger and N. Lambert, Phys. Rev. D77, 065008 (2008)
- [6] J. Bagger and N. Lambert, JHEP. 0802, 105 (2008)
- [7] A. Gustavsson, Nucl. Phys. B811, 66 (2009)
- [8] M. A. Bandres, A. E. Lipstein and J. H. Schwarz, JHEP. 0809, 027 (2008)
- [9] A. Mauri and A. C. Petkou, Phys. Lett. B666, 527 (2008)
- [10] M. Faizal, Phys. Rev. D84, 106011 (2011)
- [11] S. V. Ketov and S. Kobayashi, Phys. Rev. D83, 045003 (2011)
- [12] S. Mukhi and C. Papageorgakis, JHEP. 0805, 085 (2008)
- [13] N. Seiberg and E. Witten, JHEP. 9909, 032 (1999)
- [14] M. R. Douglas and N. A. Nekrasov, Rev. Mod. Phys. 73, 977 (2001)
- [15] S. Doplicher, K. Fredenhagen and J. E. Roberts, Commun. Math. Phys. 172, 187 (1995)
- [16] A. Connes, Non commutative geometry. Academic Press, Inc. London (1990)
- [17] G. Landi, An introduction to noncommutative spaces and their geometries. Springer-Verlag (1997).
- [18] J. Madore, S. Schraml, P. Schupp and J. Wess, Eur. Phys. J. C16, 161 (2000)
- [19] D. Brace, B. L. Cerchiai and B. Zumino, Int. J. Mod. Phys. A 1751, 205 (2002)
- [20] L. Alvarez-Gaume and M. A. Vazquez-Mozo, Nucl. Phys. B668, 293 (2003)
- [21] H. Ooguri and C. Vafa, Adv. Theor. Math. Phys. 7, 53 (2003)
- [22] N. Seiberg, JHEP. 0306, 010 (2003)
- [23] J. de Boer, P. A. Grassi and P. van Nieuwenhuizen, Phys. Lett. B574, (2003) 98 (2003)
- [24] E. Chang-Young, H. Kim and H. Nakajima, JHEP. 0804, 004 (2008)

- [25] K. Araki, T. Inami, H. Nakajima and Y. Saito, JHEP. 0601, 109 (2006)
- [26] J. S. Cook, J. Math. Phys. 47, 012304 (2006)
- [27] Y. Kobayashi and S. Sasaki, Phys. Rev. D72, 065015 (2005)
- [28] N. Berkovits and N. Seiberg, JHEP. 0307, 010 (2003)
- [29] C. S. Chu and E. Sezgin, JHEP. 9712, 001 (1997)
- [30] A. F. Ferrari, M. Gomes, J. R. Nascimento, A. Yu. Petrov and A. J. da Silva, Phys. Rev. D74, 125016, (2006)
- [31] P. S. Howe and E. Sezgin, Phys. Lett. B 394, 62 (1997)
- [32] M. Aganagic, J. Park, C. Popescu and J. H. Schwarz, Nucl. Phys. B 496, 191 (1997)
- [33] P. Pasti, D. P. Sorokin and M. Tonin, Phys. Lett. B 398, 41 (1997)
- [34] I. A. Bandos, K. Lechner, A. Nurmagambetov, P. Pasti, D. P. Sorokin and M. Tonin, Phys. Rev. Lett. 78, 4332 (1997)
- [35] P. S. Howe, E. Sezgin and P. C. West, Phys. Lett. B 399, 49 (1997)
- [36] M. Fujita, W. Li, S. Ryu and T. Takayanagi, JHEP. 0906, 066 (2009)
- [37] I. R. Klebanov and A. M. Polyakov, Phys. Lett. B550, 213 (2002)
- [38] J. H. Schwarz, JHEP. 0411, 078 (2004)
- [39] C. Ahn, H. Kim, B. H. Lee and H. S. Yang, Phys. Rev. D61, 066002 (2000)
- [40] B. Chen and J. B. Wu, JHEP. 096, 0809 (2008)
- [41] M. Benna, I. Klebanov, T. Klose and M. Smedback, JHEP. 0809, 072 (2008)